

Unified Angular Momentum of Dyons

Shalini Dangwal, P. S. Bisht and O. P. S. Negi*

2nd February 2008

Department of Physics,
Kumaun University,
S. S. J. Campus,
Almora- 263601, U.A., India

Email:shalini_dangwal@rediffmail.com
ps_bisht123@rediffmail.com
ops_negi@yahoo.co.in

Abstract

Unified quaternionic angular momentum for the fields of dyons and gravito-dyons has been developed and the commutation relations for dynamical variables are obtained in compact and consistent manner. Demonstrating the quaternion forms of unified fields of dyons (electromagnetic fields) and gravito-dyons (gravito-Heavisidian fields of linear gravity), corresponding quantum equations are reformulated in compact, simpler and manifestly covariant way.

The question of existence of monopole [1, 2, 3] has become a challenging new frontier and the object of more interest in connection with quark confinement problem of quantum chromodynamics (QCD). The eighth decades of this century witnessed a rapid development of the group theory and gauge field theory to establish the theoretical existence of monopoles and to explain their group properties and symmetries. Keeping in mind 't Hooft's solutions [4, 5] and the fact that despite of the potential importance of monopoles, the formalism necessary to describe them has been clumsy and manifestly non-covariant, a self consistent quantum field theory of generalized electromagnetic fields associated with dyons (particle carrying electric and magnetic charges) has been developed [6, 7]. The analogy between linear gravitational and electromagnetic fields leads

*Address from July 01 to August 31, 2006:- Universität Konstanz, Fachbereich Physik, Prof. Dr. H. Dehnen, Post fach M 677, D-78457 Konstanz, Germany.

the asymmetry in Einstein's linear equation of gravity and then suggests the existence of gravitational analogue of magnetic monopole [8, 9]. Like magnetic field, Cantani [10] introduced a new field (called Heavisidian field) depending upon the velocities of gravitational charges (masses) and derived the covariant equations (Maxwell's like equations) of linear gravity. Avoiding the use of arbitrary string variable [1, 2], we have also formulated manifestly covariant theory of gravito-dyons [11, 12] in terms of two four-potentials leading to the structural symmetry between generalized electromagnetic fields of dyons [13, 14] and generalised gravito-Heavisidian fields of gravito-dyons.

In this paper, we have used quaternion analysis to combine the complex description of dyons and gravito-dyons. Generalized fields of dyons and gravito-dyons are combined together and the corresponding covariant field equations and equation of motion have been derived. It has been shown that the theory leads four different Chirality's parameters associated respectively with electric, magnetic, gravitational and Heavisidian charges. The study of gauge invariant and rotationally symmetric angular momentum operator for unified quaternionic fields of dyons and gravito-dyons has been undertaken and the commutation relations associated with the components of angular momentum and other dynamical parameters have been derived. The symmetry of these commutation relations requires an additional potential term, of appropriate strength depending upon the magnetic, gravitational and Heavisidian parameter, besides the usual Colombian potential like term, in the Hamiltonian. It has been shown that this unified theory reproduces the dynamics of individual charges (masses) in the absence of others.

In order to formulate unified theory of generalized electromagnetic fields (associated with dyons) and generalized gravito-Heavisidian fields (associated with gravito-dyons) of linear gravity, we describe the properties of quaternion algebra with the use of natural units ($c = \hbar = 1$), and gravitational constant is taken unity. Unified quaternionic charge is described as [15],

$$Q = (e, g, m, h) = e - i g - j m - k h, \quad (1)$$

where e, g, m, h are respectively electric, magnetic, gravitational and Heavisidian charges (masses) and i, j, k are the quaternion units. The properties of quaternion are described as under

$$\begin{aligned} ij &= -ji = k, & i(ij) &= (ii)j = -j, \\ (ij)j &= i(jj) = -i, & ik &= -ki = -j, \\ kj &= -jk = -i, & i^2 &= j^2 = k^2 = -1. \end{aligned} \quad (2)$$

Complex structure (e, g) represents the generalized charge of dyons in electromagnetic fields while (m, h) is the generalized charge of gravito-dyons. The norm of unified quaternion charge is expressed as,

$$N(Q) = Q\bar{Q} = \bar{Q}Q = e^2 + g^2 + m^2 + h^2, \quad (3)$$

where

$$\bar{Q} = (e, -g, -m, -h) = e + i g + j m + k h. \quad (4)$$

The interaction of a^{th} quaternionic charge Q_a in the field of b^{th} quaternionic charge Q_b depends on the quantity

$$\bar{Q}_a Q_b = (W_{ab}, X_{ab}, Y_{ab}, Z_{ab}), \quad (5)$$

where

$$\begin{aligned} W_{ab} &= e_a e_b + g_a g_b + m_a m_b + h_a h_b, \\ X_{ab} &= e_a g_b - g_a e_b + m_a h_b - h_a m_b, \\ Y_{ab} &= e_a m_b - m_a e_b + h_a g_b - g_a h_b, \\ Z_{ab} &= e_a h_b - h_a e_b + g_a m_b - m_a g_b. \end{aligned} \quad (6)$$

These four coupling parameters $W_{ab}, X_{ab}, Y_{ab}, Z_{ab}$ may be identified as electric, magnetic, gravitational and Heavisidian coupling parameters associated with the basis elements $(1, i, j, k)$ of a quaternion respectively. The coupling parameter W_{ab} may also be obtained by taking the scalar product of quaternions Q_a and Q_b and thus equals to the norm given by equation (3) when $a = b$. It shows that if same quaternions were interacting with each other, their behavior would be Coulomb like. W_{ab} may then be called Coulomb like coupling parameter while the second one X_{ab} corresponds to the combined Chirality's parameters associated with the interaction of two dyons (electromagnetic and gravito-dyons). Parameter Y_{ab} corresponds to the interaction between two dyons combined by electric-gravitational and magnetic-Heavisidian charges while the fourth one Z_{ab} is the coupling parameter corresponds to the interaction of two dyons combined by the coupling of electric-Heavisidian and magnetic-gravitational charges. Unified quaternion valued potential of dyons may then be defined as,

$$V_\mu = (A_\mu, B_\mu, C_\mu, D_\mu) = A_\mu - i B_\mu - j C_\mu - k D_\mu, \quad (7)$$

or more generally we write as

$$V = (A, B, C, D) = A - i B - j C - k D, \quad (8)$$

where

$$\begin{aligned} A &= (A_0, A_1, A_2, A_3) = A_0 - i A_1 - j A_2 - k A_3, \\ B &= (B_0, B_1, B_2, B_3) = B_0 - i B_1 - j B_2 - k B_3, \\ C &= (C_0, C_1, C_2, C_3) = C_0 - i C_1 - j C_2 - k C_3, \\ D &= (D_0, D_1, D_2, D_3) = D_0 - i D_1 - j D_2 - k D_3. \end{aligned} \quad (9)$$

Similarly one may define the quaternion valued unified fields as

$$\mathfrak{S}_{\mu\nu} = F_{\mu\nu} - i \widetilde{F_{\mu\nu}} - j f_{\mu\nu} - k \widetilde{f_{\mu\nu}}, \quad (10)$$

where

$$\begin{aligned} F_{\mu\nu} &= A_{\mu,\nu} - A_{\nu,\mu} - \frac{1}{2} \varepsilon_{\mu\nu\sigma\rho} (B^{\sigma,\rho} - B^{\rho,\sigma}), \\ \widetilde{F_{\mu\nu}} &= B_{\mu,\nu} - B_{\nu,\mu} + \frac{1}{2} \varepsilon_{\mu\nu\sigma\rho} (A^{\sigma,\rho} - A^{\rho,\sigma}), \\ f_{\mu\nu} &= C_{\mu,\nu} - C_{\nu,\mu} - \frac{1}{2} \varepsilon_{\mu\nu\sigma\rho} (D^{\sigma,\rho} - D^{\rho,\sigma}), \\ \widetilde{f_{\mu\nu}} &= D_{\mu,\nu} - D_{\nu,\mu} + \frac{1}{2} \varepsilon_{\mu\nu\sigma\rho} (C^{\sigma,\rho} - C^{\rho,\sigma}) \end{aligned} \quad (11)$$

and

$$\begin{aligned} F_{\mu\nu,\nu} &= j_\mu^{(e)}, & \widetilde{F_{\mu\nu,\nu}} &= j_\mu^{(m)}, \\ f_{\mu\nu,\nu} &= j_\mu^{(G)}, & \widetilde{f_{\mu\nu,\nu}} &= j_\mu^{(H)}. \end{aligned} \quad (12)$$

the (‘tittle’) denotes the dual part of field tensor. In equations (11) the field tensors $F_{\mu\nu}$, $\widetilde{F_{\mu\nu}}$ are associated with generalized fields of dyons while those for $f_{\mu\nu}$, $\widetilde{f_{\mu\nu}}$ are associated accordingly with the generalized gravito-Heavisidian fields of gravito-dyons [16]. As such the quaternion valued current may then be defined as,

$$J_\mu = j_\mu^{(e)} - i j_\mu^{(m)} - j j_\mu^{(G)} - k j_\mu^{(H)}, \quad (13)$$

or in general

$$J = (j^{(e)}, j^{(m)}, j^{(G)}, j^{(H)}), \quad (14)$$

which is the quaternion valued expression for field equation of dyons and gravito-dyons i.e.

$$\mathfrak{S}_{\mu\nu,\nu} = J_\mu. \quad (15)$$

Lagrangian density may then be written as follows in terms of compact, simpler and consistent quaternion notation form i.e.

$$\mathcal{L} = -M - \frac{1}{8} \overline{\mathfrak{S}_{\mu\nu}} \mathfrak{S}_{\rho\sigma} \eta^{\mu\nu} \eta^{\rho\sigma} + \frac{1}{2} \overline{V_\mu} J_\nu \eta^{\mu\nu}, \quad (16)$$

where $\eta^{\mu\nu}$ is the flat space-time metric with signature -2 . Combined gauge covariant quaternion valued momentum may then be written as,

$$p \mapsto p - \frac{1}{2}(\overline{Q}V + Q\overline{V}) \implies p - eA - gB - mC - hD, \quad (17)$$

which leads to equation of motion into following compact and simpler form [17],

$$M \ddot{x}_\mu = \overline{Q} \mathfrak{S}_{\mu\nu} u^\nu, \quad (18)$$

where M is the mass of the particle, $\ddot{x}_\mu$ is the acceleration and u^ν is the four velocity of particle. The gauge invariant and rotationally symmetric angular momentum in the unified quaternionic field of dyons and gravito-dyons may be written as,

$$\vec{J} = \vec{r} \times (\vec{p} - \overline{Q} \vec{V}) + (X_{ab} + Y_{ab} + Z_{ab}) \frac{\vec{r}}{r}, \quad (19)$$

where $\overline{Q} \vec{V}$ is the quaternionic parts. The angular momentum given by equation (19) leads to following commutation relations,

$$\begin{aligned} [J_k, J_l] &= i \varepsilon_{klm} J_m \quad (i = \sqrt{-1}) \quad (k, l, m = 1, 2, 3), \\ [J^2, J_k] &= [J^2, J_l] = [J^2, J_m] = 0, \\ [J_k, r_l] &= i \varepsilon_{klm} r_m, \quad [\pi_k, r_l] = -i \delta_{kl}, \\ [\pi_k, \pi_l] &= i (X_{ab} + Y_{ab} + Z_{ab}) \varepsilon_{klm} \psi_m, \\ [J_k, \pi_l] &= i \varepsilon_{klm} \pi_m - i (X_{ab} + Y_{ab} + Z_{ab}) r_l \psi_k, \end{aligned} \quad (20)$$

where π_l is defined as,

$$\vec{\pi} = \vec{p} - \overline{Q} \vec{V}. \quad (21)$$

Keeping in mind the usual Coulombian interaction problem of electric charges, we except the following manifestly gauge invariant and rotationally symmetric Hamiltonian for the system of unified charges i.e.

$$\mathcal{H} = \frac{\pi^2}{2m} - \frac{W_{ab}}{r} + V(r), \quad (22)$$

where $\vec{\pi}$ is the gauge invariant linear momentum described by equation (21). The additional potential term $V(r)$ may be identified and decided according to the symmetry requirements of the system. Using equation (19), the value of the operator J^2 may be calculated as

$$J^2 = (\vec{r} \times \vec{\pi}) \cdot (\vec{r} \times \vec{\pi}) + (X_{ab}^2 + Y_{ab}^2 + Z_{ab}^2) + 2(X_{ab} + Y_{ab} + Z_{ab}) \vec{r} \cdot (\vec{r} \times \vec{\pi}) \quad (23)$$

The third term in equation (23) is zero and hence we may write,

$$\frac{\pi^2}{2m} = \frac{J^2}{2mr^2} - \frac{(X_{ab}^2 + Y_{ab}^2 + Z_{ab}^2)}{2mr^2} + \frac{p^2}{2m}. \quad (24)$$

It is quite obvious that the Hamiltonian given by equation (22) possesses the higher symmetry same as that of the pure Coulomb Hamiltonian provided the additional potential $V(r)$ in equation (22) takes the scalar form,

$$V(r) = \frac{(X_{ab}^2 + Y_{ab}^2 + Z_{ab}^2)}{2mr^2}. \quad (25)$$

Thus the Hamiltonian given by equation (22) may be written as,

$$\mathcal{H} = \frac{\pi^2}{2m} - \frac{W_{ab}}{r} + \frac{(X_{ab}^2 + Y_{ab}^2 + Z_{ab}^2)}{2mr^2}, \quad (26)$$

which leads to the following commutation relations,

$$[J^2, \mathcal{H}] = [J, \mathcal{H}] = 0. \quad (27)$$

Quaternion valued unified vector field $\vec{\psi}$ is written as,

$$\vec{\psi} = (\vec{E}, \vec{M}, \vec{G}, \vec{H}), \quad (28)$$

where $\vec{E}, \vec{M}, \vec{G}, \vec{H}$ are generalized electric, magnetic, gravitational and Heavisidian fields respectively defined [15] as,

$$\begin{aligned} \vec{E} &= -\frac{\partial \vec{A}}{\partial t} - \vec{\nabla} A_0 - \vec{\nabla} \times \vec{B}, \\ \vec{M} &= -\frac{\partial \vec{B}}{\partial t} - \vec{\nabla} B_0 + \vec{\nabla} \times \vec{A}, \\ \vec{G} &= -\frac{\partial \vec{C}}{\partial t} - \vec{\nabla} C_0 - \vec{\nabla} \times \vec{D}, \\ \vec{H} &= -\frac{\partial \vec{D}}{\partial t} - \vec{\nabla} D_0 + \vec{\nabla} \times \vec{C}. \end{aligned} \quad (29)$$

These generalized electric, magnetic, gravitational and Heavisidian fields satisfy the following pair of Generalized Dirac Maxwell's (GDM) equations of dyons and gravito-dyons [11, 12];

$$\vec{\nabla} \cdot \vec{E} = j_0^{(e)}, \quad \vec{\nabla} \cdot \vec{M} = j_0^{(m)},$$

$$\begin{aligned}
\vec{\nabla} \cdot \vec{G} &= j_0^{(G)}, \quad \vec{\nabla} \cdot \vec{H} = j_0^{(H)}, \\
\vec{\nabla} \times \vec{E} &= -\frac{\partial \vec{M}}{\partial t} - \vec{j}^{(m)}, \quad \vec{\nabla} \times \vec{M} = \frac{\partial \vec{E}}{\partial t} + \vec{j}^{(e)}, \\
\vec{\nabla} \times \vec{G} &= \frac{\partial \vec{H}}{\partial t} + \vec{j}^{(H)}, \quad \vec{\nabla} \times \vec{H} = -\frac{\partial \vec{G}}{\partial t} - \vec{j}^{(G)},
\end{aligned} \tag{30}$$

where $(j_0^{(e)}, j_0^{(m)}, j_0^{(G)}, j_0^{(H)})$ are respectively charge densities for electric, magnetic, gravitational and Heavisidian charges while $(\vec{j}^{(e)}, \vec{j}^{(m)}, \vec{j}^{(G)}, \vec{j}^{(H)})$ are corresponding current source densities due to these charges. Quaternion unified potential V given by equation (7) and unified field $\vec{\psi}$ are then related as,

$$\vec{\psi} = -\frac{\partial \vec{V}}{\partial t} - \vec{\nabla} V_0 + \vec{\nabla} \times \vec{V}. \tag{31}$$

Operating the quaternionic differential operator $\partial = \partial_0 - i\partial_1 - j\partial_2 - k\partial_3$ to equation (7), we get the following unified form of quaternionic potential equation,

$$\partial V = \vec{\psi}, \tag{32}$$

which is quaternion valued unified potential equation for generalized charges of dyons. Unified quaternion valued current may then be written as,

$$J = (j^{(e)}, j^{(m)}, j^{(G)}, j^{(H)}), \tag{33}$$

where

$$\begin{aligned}
\partial \bar{\partial} A &= \square A = j^{(e)}, \\
\partial \bar{\partial} B &= \square B = j^{(m)}, \\
\partial \bar{\partial} C &= \square C = j^{(G)}, \\
\partial \bar{\partial} D &= \square D = j^{(H)}, \\
\partial \bar{\partial} &= \square = \partial_\mu \partial_\mu = -\frac{\partial^2}{\partial t^2} + \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}.
\end{aligned} \tag{34}$$

Quaternion field equation associated with unified quaternion charge may then be written as,

$$\partial \bar{\partial} V = \square V = J. \tag{35}$$

Quaternion valued field tensor density is accordingly expressed as

$$Q_{\mu\nu} = (A_{\mu\nu}, B_{\mu\nu}, C_{\mu\nu}, D_{\mu\nu}), \tag{36}$$

where

$$\begin{aligned}
Q_{\mu\nu} &= V_{\mu,\nu} - V_{\nu,\mu}, \\
A_{\mu\nu} &= A_{\mu,\nu} - A_{\nu,\mu}, \\
B_{\mu\nu} &= B_{\mu,\nu} - B_{\nu,\mu}, \\
C_{\mu\nu} &= C_{\mu,\nu} - C_{\nu,\mu}, \\
D_{\mu\nu} &= D_{\mu,\nu} - D_{\nu,\mu},
\end{aligned} \tag{37}$$

are respectively field tensors of electric, magnetic, gravitational and Heavisidian charges and comma (,) denotes partial differentiation. A_μ and B_μ are dual invariant for generalized electromagnetic fields of dyons while C_μ and D_μ are dual invariant under duality transformations for generalized fields of gravito-dyons. Unified quaternion valued current J_μ and field tensor $Q_{\mu\nu}$ are then related in the following manner;

$$Q_{\mu\nu,\nu} = J_\mu, \tag{38}$$

which is equivalent to the following quaternionic form;

$$\bar{\partial}\psi = J. \tag{39}$$

as such, an unified field theory of generalised electromagnetic and Heavisidian fields have been developed in view of the fact that these fields essentially possess the structural symmetry. Quaternion charge defined by equation (1) represents the theory of particles carrying simultaneously electric, magnetic, gravitational and Heavisidian charges. Equation (1) has been described as the combination of two complex charges of dyons (e, g) and gravito-dyons (m, h) . Equation (6) represents the coupling parameters W_{ab} , X_{ab} , Y_{ab} and Z_{ab} . This equation shows that in the presence of only electric charge the coupling becomes $W_{ab} = e_a e_b$ while the other parameters X_{ab}, Y_{ab} and Z_{ab} vanish. If the particles are considered as dyons (e, g) of electromagnetic fields, we have $W_{ab} = e_a e_b + g_a g_b$ and $X_{ab} = e_a g_b - g_a e_b$ which are respectively named as electric and magnetic coupling parameters. For gravito-dyons (m, h) we have only $W_{ab} = m_a m_b + h_a h_b$ and $X_{ab} = m_a h_b - h_a m_b$. For other dyon like (e, h) i.e. electric and Heavisidian charges we get $W_{ab} = e_a e_b + h_a h_b$ and $Z_{ab} = e_a h_b - h_a e_b$. For other dyon like (g, m) i.e. gravitational charge and magnetic monopole, we get $W_{ab} = m_a m_b + g_a g_b$ and $Z_{ab} = g_a m_b - m_a g_b$. For other dyon like (e, m) i.e. electric charge and gravitational charge (mass), we get $W_{ab} = e_a e_b + m_a m_b$ and $Y_{ab} = e_a m_b - m_a e_b$. Similarly for purely hypothetical dyons (h, g) i.e. Heavisidian and magnetic charges we get $W_{ab} = h_a h_b + g_a g_b$ and $Y_{ab} = h_a m_b - m_a h_b$. As such, the unified theory of quaternionic charge is the combined theory of fields characterizing the study of fields associated with various types of dyons.

Equation (7) represents unified quaternion valued potential of dyons. It is to be noted that here we have combined four-charges (electric, magnetic,

gravitational and Heavisidian) in terms of tetrad combination in the context of special relativity (not in terms of generalised coordinates) and maintained the abelian gauge structure in terms of four photons with a the group is defined as $\mathcal{G} = \text{U}^{(e)}(1) \times \text{U}^{(m)}(1) \times \text{U}^{(G)}(1) \times \text{U}^{(H)}(1)$. While the quaternion structure leads to rotation in charge space and it is our next target to find out these investigations of quaternion rotation shortly to mix up the gauge structures. here we have the possibility to consider the gravitational charge m both positive or negative. This should imply that gravitational and Heavisidian charges may be either positive or negative dealing with the concept of negative masses. On the other hand, properties of quaternion algebra are incorporated to derive the field equation (10) as a quaternion and the coefficients of its basis elements represent the algebraic gauge structure field equations. Lagrangian density (16) has been shown to yield the compact and simpler representation of combined dyonic field equation (15) and equation of motion (18). The gauge invariant and rotationally symmetric angular momentum in the unified quaternionic fields of dyons and gravito-dyons given by equation (19) contains an extra term $J_{res} = (X_{ab} + Y_{ab} + Z_{ab}) \frac{\vec{r}}{r}$ of residual angular momentum in addition to the usual kinetic angular momentum $J_{kin} = \vec{r} \times (\vec{p} - \vec{Q} \vec{V})$. The rotationally symmetric nature of angular momentum is also reflected in the commutation relation given by equation (20). The commutation relations given by equations (20) and (27) show that the operator and are the constant of motion. The commutation relation (27) demands an additional term in the Hamiltonian so that it consists the higher symmetry as the pure Coulomb Hamiltonian. From the above analysis it may be concluded that besides the potential importance of monopole as intrinsic part of grand unified theories, monopoles and dyons may provide even more ambitious model to purport the unification of gravitation with strong and electro-weak forces. The electric, magnetic, gravitational and Heavisidian fields defined by equation (29) describe the pair of Maxwell's equations associated with generalized fields of dyons (e, g) and gravito-dyons (m, h) . Equation (39) describes the unified forms of quaternion four-current density and it may be shown that these quaternion field equations are invariant under quaternion and duality transformations.

Acknowledgment: - O. P. S. Negi is thankful to Prof. Dr. H. Dehnen, Universität Konstanz, Fachbereich Physik, Postfach M 677, D-78457 Konstanz, Germany for his hospitality at Konstanz. He is also grateful to German Academic Exchange Service (DAAD) for financial support under re-invitation fellowship programme.

References

- [1] P. A. M. Dirac, Proc. Roy. Soc. London, **A133**, 60 (1931).
- [2] P. A. M. Dirac, Phys. Rev., **74**, 817 (1948).
- [3] J. Schwinger, Science, **165**, 757 (1978).

- [4] G. 't Hooft, Nucl. Phys., **B79** , 276 (1974).
- [5] G. 't Hooft, Nucl. Phys., **B138** , 1 (1978).
- [6] B. S. Rajput and D. C. Joshi, Hadronic J., **4** , 1805 (1981).
- [7] B. S. Rajput and Om Prakash, Indian J. Phys., **A53** , 274 (1979).
- [8] J. S. Dowker and J. A. Roche, Proc. R. Soc., **92** , 1 (1967).
- [9] A. Singh, Lett. Nuovo Cimento, **32** , 232 (1981).
- [10] D. D. Cantani, Nuovo Cimento, **A60** , 67 (1980).
- [11] P. S. Bisht, O. P. S. Negi and B. S. Rajput, Ind. J. P. and App. Phys., **28** , 157 (1990).
- [12] P. S. Bisht, O. P. S. Negi and B. S. Rajput, Prog. Theor. Phys., **85** , 157 (1991).
- [13] D. S. Bhakuni, O. P. S. Negi and B. S. Rajput, Lett. Nuovo Cimento, **36** , 499 (1983).
- [14] B. S. Rajput, S. R. Kumar and O. P. S. Negi, Lett. Nuovo Cimento, **36** , 75 (1983).
- [15] P. S. Bisht, O. P. S. Negi and B. S. Rajput, Ind. J. P. and Appl. Maths., **24** , 543 (1993).
- [16] P. S. Bisht, O. P. S. Negi and B. S. Rajput, Int. J. Theor. Phys., **32** , 2099(1993).
- [17] P. S. Bisht, O. P. S. Negi and B. S. Rajput, IL Nuovo Cimento, **A 104** , 337 (1991).